




Scalable geospatial data generation with AlphaEarth

NeurIPS 2025

Tackling Climate Change with Machine Learning




Motivation

Global damage costs from natural disasters, Wildfire, 1980 to 2024




Total economic cost of damages as a result of global natural disasters in any given year, measured in current US\$.

Damage Costs from Wildfires (Bn USD)




Source: Our World in Data, EM-DAT, CREED / UCLouvain (2024)
Note: Data includes disasters recorded up to April 2024





Wildfires know no borders...















... But high quality vegetation datasets do!


- LANDFIRE is an American crossagency program
- Produces vegetation dataset in the US for fire management
- EVT: "Existing Vegetation Type"





"Landfire technical documentation." (2023) Washington DC: US Department of the Interior, US Geological Survey


About AlphaEarth Foundations (AEF)



- Google Deepmind model-as-data publicly released on Google Earth Engine
- Model trained on wide variety of geospatial data (satellite optical + SAR, ERA5...)
- 'General purpose' embeddings provided as 64-dimensioned vectors


Brown, Christopher F., et al. "AlphaEarth Foundations: An embedding field model for accurate and efficient global mapping from sparse label data." arXiv preprint arXiv:2507.22291 (2025).

Project overview



Inference in Canada


13 classes





Methods

• 2 held out test areas to evaluate performance

• 4 different models investigated







Training and testing

- Similar performance across models
- Drop in performance (a) in test regions and (b) as number of classes increases
- Over-performing random forest model in training

13 classes

	ACC	J	F 1	ACC	J	F 1	ACC	J	F 1
Logistic Regression	0.77	0.48	0.60	0.77	0.48	0.59	0.71	0.39	0.51
Random Forest	0.97	0.95	0.97	0.81	0.55	0.67	0.73	0.43	0.55
Gradient Boosted Trees	0.79	0.52	0.65	0.79	0.52	0.64	0.73	0.42	0.54
Segmentation Model	0.79	0.50	0.63	0.79	0.51	0.63	0.73	0.42	0.54

Validation

Test

Training

80 Classes


	Training		Validation			Test			
	ACC	J	F 1	ACC	J	F 1	ACC	J	F 1
Logistic Regression	0.60	0.28	0.40	0.60	0.27	0.39	0.42	0.11	0.17
Random Forest	0.96	0.94	0.96	0.71	0.39	0.53	0.48	0.16	0.23
Gradient Boosted Trees	0.63	0.29	0.43	0.62	0.28	0.41	0.44	0.12	0.17
Segmentation Model	0.65	0.28	0.40	0.66	0.29	0.41	0.48	0.15	0.21


Training and testing


 Large variations in performances across test regions

13 classes

	Abrunt transition in LANDEIDE	dota could ovalain	Canada South			Canada West			
	Abrupt transition in LANDFIRE drop in performance	data could explain	ACC	J	F 1	ACC	J	F1	
		Logistic Regression	0.67	0.31	0.41	0.82	0.35	0.42	
		Random Forest	0.69	0.34	0.45	0.83	0.42	0.52	
		Gradient Boosted Trees	0.69	0.34	0.45	0.83	0.32	0.39	
		Segmentation Model	0.69	0.34	0.45	0.83	0.37	0.45	









- EVT is itself the output of a predicted model not a 'true ground truth'
- Performance drops as a function of distance to labeled data
- Simple models can be used for this task with AlphaEarth Foundations
- This study can be extended to other use cases outside of vegetation types