



# Harrell's Doveweed Postemergence Control Guide

## Turf Safety

Product	HRAC Code	Creeping Bentgrass	Fine Fescue	Kentucky Bluegrass	Perennial Ryegrass	Rough Bluegrass	Tall Fescue	Bahiagrass	Bermudagrass	Buffalograss	Carpetgrass	Centipedegrass	Kikuyugrass	Seashore Paspalum	St. Augustinegrass	Zoysiagrass
Atrazine	5	-	-	-	-	-	-	-	L	-	-	L	-	-	L	L
Blindside	14 + 2	-	-	L	-	-	L	-	L	L	-	L	-	-	L	L
Celsius	2 + 4	-	-	-	-	-	-	-	L	L	-	L	-	-	L	L
GameOn	4	L	L	L	L	-	L	-	L	-	-	-	-	-	-	L
Simazine	5	-	-	-	-	-	-	-	L	-	-	L	-	-	L	L
Speedzone	4 + 14	L	L	L	L	-	L	-	L	L	-	-	-	-	-	L
Surge	4 + 14	L	L	L	L	-	L	L	L	L	-	L	-	-	-	L
<b>Tribute Total</b>	<b>2</b>	-	-	-	-	-	-	-	L	-	-	-	-	-	-	L

\*L = Labeled; - = Not Labeled. Consult product label for specific turf safety limitations.

## Use Sites

Product	Residential	Commercial	Golf Course	Sod Farms	Sports Fields
Atrazine	X	-	X	X	X
Blindside	X	X	X	X	X
Celsius	X	X	X	X	X
GameOn		X	X	X	
Simazine	X	X	X	X	X
Speedzone	X	X	X	X	X
Surge	X	X	X	X	X
<b>Tribute Total</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>

\*X = Labeled use site. Consult product label for specific use site limitations.

# Harrell's Doveweed Postemergence Control Guide

## Application Programs

Product	Program*
<b>Atrazine</b>	Apply 1 lb active ingredient/A. Apply a sequential application a minimum 30 days after initial application or once regrowth is observed, whichever is longer. Application on a granular carrier will limit phytotoxicity. Irrigate following application. Treatment will provide both post- and preemergence control.
<b>Blindside</b>	Apply 5 oz/A followed by a sequential application 14-21 days after the initial. For long lasting control, apply in combination with a pre-emergent herbicide.
<b>Celsius</b>	Apply at 4.9 oz/A. Apply a sequential application 2-4 weeks after initial application or after regrowth is observed. Do not exceed 7.4 oz/A per year for all applications. Add Harrell's Nonionic Penetrant Plus at 0.25% v/v to the spray solution. For long lasting control, apply in combination with a pre-emergent herbicide.
<b>GameOn</b>	Apply 3 pt/A. Add Harrell's Nonionic Penetrant Plus at 0.25% v/v to the spray solution. Apply a sequential application 3 weeks after initial application or after regrowth is observed, whichever is longer. For long lasting control, apply in combination with a pre-emergent herbicide.
<b>Simazine</b>	Apply 2 lb active ingredient/A. Apply a sequential application at least 30 days after the initial application, but not after June 1. Do not exceed 3 lb active ingredient/A per calendar year. Irrigate following application. Treatment will provide both post- and preemergence control.
<b>Speedzone</b>	Apply at 4 pt/A. Apply a sequential application a minimum 30 days after initial application or once regrowth is observed, whichever is longer. For long lasting control, apply in combination with a pre-emergent herbicide.
<b>Surge</b>	Apply 3.25 pt/A. Apply a sequential application a minimum 30 days after initial application or once regrowth is observed, whichever is longer. For long lasting control, apply in combination with a pre-emergent herbicide.
<b>Tribute Total</b>	Apply 3.2 oz/A. Include Harrell's Methylated Seed Oil at 0.5-1% v/v and 1.5-3 lb/A ammonium sulfate in the spray solution. Apply late July/early August. Make a repeat application at regrowth. For long lasting control, apply in combination with a pre-emergent herbicide such as Specticle.

\*Consult product label for specific use instructions and product restrictions.

