

SECUKINUMAB (Cosentyx) Infusion Order Form

		Skownegan, Maine 04976
Diagnosis [] Psoriatic arthritis [] Other:		
Weight:kg Height:inches] New treatment] Transfer – Receiving secukinumab since
Negative TB test date	L	
 Prior to initial therapy: (Provide copies if results not repo Pregnancy test if between 14 and 50 years old with g Hepatits C virus antibody, HIV, C-reactive protein Screen for Hepatitis B infection: Hep B surface antig TB screen with PPD or IGRA before start of therapy 	estationa en, Hep	B Antibody, Hep B Core Antibody total
PRE-MEDICATE (30 minutes before infusion) or		
[] acetaminophen 1000 mg PO		[] Methylprednisolone 40 mg IV
[] loratadine 10 mg PO		[] Hydrocortisone 50 mg IV
[] diphenhydramine 25 mg PO		[] Other:
Secukinumab IV		[] Outer.
[] Initial – with loading dose - 6 mg/kg at week Then 1.75 mg/mg (max 300 mg) every		[] Maintenance or omit loading dose 1.75 mg/mg (max 300 mg) every 4 weeks
Instruct patient to seek medical attention if they [X] Repeat TB screen with IGRA if no IGRA or F [] CBC and hepatic panel - recommended at 3 to [] Other:	aintenanter. infusio feel sic PPD in t 6 montl	on. May be discharged if post-infusion vital signs are stable. ek or develop a fever, cough, or other signs of infection. the past 12 months hs after initiating therapy. [] pending [] complete [] not needed# at health insurer who authorized.
		oraumorization:
Checklist for non-RFGH credentialed providers: Annually: [] H&P completed with last year If new therapy - attach:		
FAX to RFGH Infusion clinic at 207-858-2404		ontact Infusion Clinic at 207-858-8722
Provider signature		Prep: 9/2
Printed name		
	Pa	tient

dob

phone #